

Pion-photon transition form factor with analytic coupling

I. A. Zemlyakov^{†*}, I. L. Chuev[×], A. V. Kotikov^{°1)}

[†]Department of Physics, Universidad Tecnica Federico Santa Maria,
1680 Avenida Espana, Valparaiso, Chile

^{*}Tomsk State University, 634010 Tomsk, Russia

[×]Department of Theoretical Physics, Moscow Institute for Physics and Technology,
141701 Dolgoprudny, Russia

[°]Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Russia

Submitted 1 August 2026

Resubmitted 28 August 2026

Accepted 29 August 2026

We investigate the pion-photon transition form factor within the framework of analytic QCD. A comparison is performed between experimental data and perturbative QCD based on conventional and analytic versions of perturbation theory with the “massive” form of the twist-four contribution. We show that conventional perturbation theory fails to reproduce the data, while the analytic version demonstrates good agreement with experiment.

DOI: 10.7868/S3034576626100013

The results obtained in analytic QCD are very stable with increasing order of perturbation theory. Moreover, they are in perfect agreement with the results in [1, 2] (see Fig. 1), obtained in the framework of so-called method of lightcone sum rules in forms of dispersion relations [3, 4].




Fig. 1. (Color online) The results in the next-to-next-to-leading order of analytic QCD together with the results obtained in [1]

Funding. One of us (I. A. Zemlyakov) was supported in part by the Fellowship ANID Beca de Doctorado Nacional # 21250067. A. V. Kotikov was supported by the Russian Science Foundation grant # 25-22-00576.

Conflict of interest. The authors of this work declare that they have no conflicts of interest.

1. S. V. Mikhailov, A. V. Pimikov, and N. G. Stefanis, “Extending the application of the light-cone sum rules method to low momenta using QCD renormalization-group summation: Theory and phenomenology”, *Phys. Rev. D* **103**(9), 096003 (2021).
2. S. Mikhailov, A. Pimikov, and N. G. Stefanis, “Improved calculation of the $\gamma^*\gamma \rightarrow \pi$ process at low Q_2 using LCSR’s and renormalization-group summation”, *EPJ Web Conf.* **258**, 03003 (2022).
3. I. I. Balitsky, V. M. Braun, and A. V. Kolesnichenko, “Radiative Decay $\Sigma^+ \rightarrow p\gamma$ in Quantum Chromodynamics”, *Nucl. Phys. B* **312**, 509 (1989).
4. A. Khodjamirian, “Form-factors of $\gamma^*\rho \rightarrow \pi$ and $\gamma^*\gamma \rightarrow \pi^0$ transitions and light cone sum rules”, *Eur. Phys. J. C* **6**, 477 (1999).

¹⁾e-mail: kotikov@theor.jinr.ru