

**П И С Ь М А**  
**В ЖУРНАЛ ЭКСПЕРИМЕНТАЛЬНОЙ**  
**И ТЕОРЕТИЧЕСКОЙ ФИЗИКИ**

ОСНОВАН В 1965 ГОДУ  
ВЫХОДИТ 24 РАЗА В ГОД

ТОМ 71, ВЫПУСК 4  
25 ФЕВРАЛЯ, 2000

Pis'ma v ZhETF, vol.71, iss.4, pp.187 - 191

© 2000 February 25

**TEST OF THE QCD VACUUM WITH THE SOURCES IN HIGHER  
REPRESENTATIONS**

Yu.A.Simonov

*State Research Center Institute of Theoretical and Experimental Physics  
117259 Moscow, Russia*

Submitted 20 December 1999

Resubmitted 20 January 2000

Recent accurate measurement [1, 2] of static potentials between sources in various  $SU(3)$  representations provides a crucial test of the QCD vacuum and of different theoretical approaches to the confinement. In particular, the Casimir scaling of static potentials found for all measured distances implies a strong suppression of higher cumulants and a high accuracy of the Gaussian stochastic vacuum. Most popular models are in conflict with these measurements.

PACS: 12.38.-t

1. An accurate measurement of static potentials between sources in the eight different representations of  $SU(3)$  group made recently in [1] reveals a new quantitative picture of the QCD vacuum and provides a crucial test of existing theoretical models. Other measurements of static interaction [2] are in general agreement with [1].

The most useful way to represent static potentials  $V_D(r)$  in representations  $D = 3, 8, 6, 15a, 10, 27, 24, 15s$  is through the complete set of field correlators in the framework of the Field Correlator Method (FCM) [3]:

$$V_D(r) = - \lim_{T \rightarrow \infty} \frac{1}{T} \ln \langle W(C) \rangle, \quad (1)$$

where Wilson loop  $W(C)$  for the rectangular contour  $C = r \times T$  in the (34) plane has the cumulant expansion,

$$\langle W(C) \rangle = Tr_D \exp \int_S \sum_{n=2,4,\dots} (ig)^n \langle \langle F(1)F(2)\dots F(n) \rangle \rangle d\sigma(1)\dots d\sigma(n). \quad (2)$$

Here  $F(k)d\sigma(k) = F_{34}(u^{(k)}, x_0)d\sigma_{34}^{(k)}$  and the component  $F_{34}(u, x_0) \equiv E_3(u, x_0) = \phi(x_0, u)E_3(u)\phi(u, x_0)$ , where  $\phi$  is a parallel transporter and  $x_0$  is an arbitrary point on the surface  $S$  inside contour  $C$ ;  $Tr_D \hat{1} = 1$ .

© 2000 Российская академия наук, Отделение общей физики и астрономии,  
Институт физических проблем им. П.Л.Капицы.

Dependence on  $D$  enters in (2) through the generators  $T^a$ , since  $F(k) = F^a(k)T^a$  ( $a = 1, \dots, N_c^2 - 1$ ), and the main characteristics of  $D$  is the quadratic Casimir operator  $C_D$ :  $T^a T^a = \hat{1} C_D$ , so that the invariant square of the color charge in the representation  $D$  is  $g^2 C_D$ .

One can now express the connected correlators (cumulants) in (2) via  $C_D$  and  $D$ -independent averages as follows (for more details see last reference in [3] and [4]),

$$\text{Tr}_D \langle F(1)F(2) \rangle = C_D \frac{\langle F^a(1)F^a(2) \rangle}{N_c^2 - 1}, \quad (3)$$

$$\begin{aligned} \text{Tr}_D \langle \langle F(1)F(2)F(3)F(4) \rangle \rangle &= \frac{C_D^2}{(N_c^2 - 1)^2} \{ \langle F^a(1)F^a(2)F^b(3)F^b(4) \rangle \} + \\ &+ \langle F^a(1)F^b(2)F^b(3)F^a(4) \rangle - \langle F^a(1)F^a(2) \rangle \langle F^b(3)F^b(4) \rangle + \\ &+ (1 - \frac{N_c}{2C_D}) \langle F^a(1)F^b(2)F^a(3)F^b(4) \rangle \} + O(\frac{1}{N_c^2}). \end{aligned} \quad (4)$$

Note that the arguments of  $F(k)$  in (4) and in (2) are ordered (e.g. clockwise,  $u^{(1)} < u^{(2)} < u^{(3)} < u^{(4)}$ ) and therefore the only vacuum insertion is possible in the first term on the r.h.s. of (4) leading to the cancellation with the third term: hence the correlator (4) is a connected one vanishing at large distances,  $|u^{(1)} + u^{(2)} - u^{(3)} - u^{(4)}| \rightarrow \infty$ .

One can show in a similar way that the  $n$ -th cumulant in (2) contributes proportionally to  $C_D^n$ . As a result the static potential  $V_D(r)$  has the expansion

$$V_D(r) = d_D V^{(2)}(r) + d_D^2 V^{(4)}(r) + \dots, \quad (5)$$

where in notations of ref. [1]  $d_D = C_D/C_F$  and  $C_F$  is the fundamental Casimir operator,  $C_F = (N_c^2 - 1)/2N_c$ . The fundamental static potential contains perturbative Coulomb part  $V_{Coul}$ , confining linear and constant terms.

The Coulomb part, which is obtainable from the perturbative contribution to FC, is now known up to two loops [5] and is proportional to  $C_D$ . Therefore one may expect quartic contributions proportional to  $C_D^2 \sim d_D^2$ , to the constant and linear terms, writing (5) as

$$V_D(r) = d_D V^{(2)}(r) + d_D^2 (\bar{v}_0^{(4)} + \bar{\sigma}_4 r). \quad (6)$$

Here  $\bar{v}_0, \bar{\sigma}_4$  measure the contribution of the quartic cumulants to the constant term and string tension respectively.

Now the measurements of ref. [1] allow to find  $\bar{v}_0, \bar{\sigma}_4$  from all 8 sets of data. To this end one forms 7 combinations  $\zeta_D \equiv V_D(r) - d_D V_F(r) = d_D(d_D - 1)(\bar{v}_0^{(4)} + \bar{\sigma}_4 r)$ . As a typical example - take fundamental and adjoint potentials at distances  $r$  in the interval between 0.05 fm and 1.1 fm from the data [1] and the  $\chi^2$  fit yields for  $\bar{v}_0, \bar{\sigma}_4$ :

$$\bar{v}_0^{(4)} = (-0.6 \pm 0.67) \cdot 10^{-3} \text{ GeV}, \quad (7)$$

$$\bar{\sigma}_4 = (-2 \pm 0.69) \cdot 10^{-3} \text{ GeV}^2. \quad (8)$$

The quality of the fit is reasonable,  $\chi^2/N = 0.45$ ,  $N = 43$ . One obtains similar figures for  $D = 6, 15a, 10$  (while 3 higher representations do not yield additional information) suggesting that  $\bar{\sigma}_4^{(4)}$  is negative, and  $\bar{v}_0^{(4)}$  is compatible with zero, confirming in this way the parametrization (6).

This analysis demonstrates the phenomenon of the Casimir scaling, i.e. proportionality of static potential  $V_D(r)$  to the Casimir operator  $C_D$  with accuracy better than one percent.

Physical consequences of the Casimir scaling are numerous and important.

First of all, the sign and magnitude of quartic correction (7),(8) can be understood in the FCM. Indeed the quartic term enters the potential  $V_D$  with the factor  $(-g^4)$ , as compared to  $+g^2$  for the quadratic (Gaussian) term. Secondly, - estimate  $\langle E_3^2 \rangle$  term from the standard gluonic condensate as follows:

$$g^2 \langle E_3^a E_3^a \rangle \sim \frac{4\pi^2}{12} (0.04 \pm 0.02) \text{ GeV}^4 \sim (0.10 \pm 0.06) \text{ GeV}^4 \quad (9)$$

and take into account that the cumulant expansion in (2) is actually in powers of the parameter

$$\xi \equiv g^2 \langle E_3^a E_3^a \rangle T_g^4. \quad (10)$$

Here  $T_g$  is the correlation length of the QCD vacuum; for bilocal correlator it was measured on the lattice [6],  $T_g^{(2)} \sim 1 \text{ GeV}^{-1}$ . With the use of (10) one could expect that  $\bar{\sigma}_4$  would be from 4 to 10% of the standard string tension,  $\sigma = 0.2 \text{ GeV}^2$  provided  $T_g = T_g^{(2)}$ . The value of  $\bar{\sigma}^{(4)}$  calculated in (8) is at least 6 times smaller and suggests that quartic correlation length  $T_g^{(4)}$  may be smaller than the Gaussian one,  $T_g^{(2)} \sim 0.2 \text{ fm}$ .

This result means that the Gaussian Stochastic Model (GSM), suggested in [3] and successfully used heretofore in many applications [7], can be more accurate than it was even expected, at least in processes where string tension plays the most important role. On the other hand, the smallness of quartic and higher contributions implies a very specific picture of vacuum correlations.

Indeed the smallness of  $T_g^{(4)}$  implies that color fields tend to form compact white bilocal combinations  $F^a(1)F^a(2)$  which are almost noninteracting between themselves and therefore not contributing to the higher connected correlators. This looks like the picture of small white dipoles made of fields  $FF$  (or of vector potentials  $A_\mu A_\mu$  connected to  $FF$  in the Fock - Schwinger or contour gauge).

One can also understand qualitatively the difference between  $T_g^{(2)}$  and  $T_g^{(4)}$ , since  $T_g^{(2)}$  measures correlation length between adjoint fields  $E_3^a(x)$  and  $E_3^a(y)$  in the Gaussian correlator  $\langle E_3^a(x)E_3^a(y) \rangle$ , while  $T_g^{(4)}$  refers to the correlation of two white complexes, and should be connected to the lowest glueball mass,  $M_G \approx 2 \text{ GeV}$ ; hence  $T_g^{(4)} \sim 1/M_G \sim 0.1 \text{ fm} < T_g^{(2)}$ .

Finally, the Casimir scaling imposes severe restrictions on existing models of the QCD vacuum. For example, the center-symmetry flux model was tested and ruled out in [1], since in the original formulation it predicts vanishing adjoint string tension, while in the later modification - the fat vertex model [9] it is still far from the accurate data [1].

Next one should mention models of the abelian projected vacuum which fail to provide Casimir scaling [11], at least in the simplest version [12].

Consider now the dilute instanton gas model and the  $SU(2)$  group. The instantons may be present in the confining vacuum as an important source of chiral symmetry breaking. Then the Casimir scaling [1] imposes a strict bound on the admixture of instantons in the QCD vacuum. Indeed, insertion in (2) and (4) of the instanton field strength  $gF_{34}^a(x, z) = 4\delta_{a3}\rho^2/[(x-z)^2 + \rho^2]^2$ , ( $\rho$ -instanton size,  $z$ -its position, contribution of

parallel transporters is neglected for simplicity since it gives for bilocal correlators a reduction of 20-30%, see [11] for details) yields the following expression for the quartic contribution to the potential generated by instantons at small distances,  $r \leq \rho$  (in  $SU(2)$  group);

$$V_D^{(4)} = -\frac{N}{V} \frac{r^4}{\rho} \frac{\pi^6}{320} \frac{3C_D^2 - C_D}{15} \quad (11)$$

where  $N/V$  is the density of instanton in the vacuum. With the limit (8) on  $\bar{v}_0^{(4)}$  and taking the distance  $r = \rho = 0.3 \text{ fm}$  one obtains an upper limit on density of instantons:

$$N/V \leq 0.2 \text{ fm}^{-4} \quad (12)$$

which is much smaller than normal instanton density of  $1 \text{ fm}^{-4}$ .

With such density the role of instantons in chiral symmetry breaking and other effects would be rather small. Even more stringent bounds can be obtained from the quartic string tension, however the nonzero value of  $\bar{\sigma}_4$  for instantons does not imply confinement. One should take into account that at large distances the sum of all partial string tensions  $\sum_{n=2,4,\dots} \bar{\sigma}^{(n)}$  for the dilute instanton gas vanishes [12, 13].

In previous discussion we have ignored the fact that at large enough distances the adjoint charges are screened by the vacuum gluons, and the limiting value of the adjoint potential is equal to the doubled gluelump mass,  $2M_{gl}$ . This leads to an estimate of the screening distance  $r_0$  from the relation;  $V_{adj}(r_0) = 2M_{gl}$  where  $M_{gl}$  from [4] is around 1.4 GeV and therefore  $r_0 \approx 1.4 \text{ fm}$ , which is beyond the distance where Casimir scaling was measured in [1].

Thus the Casimir scaling is a stringent test for all models considered and displays a strong suppression of quartic and higher correlators and hence supports a good accuracy of the GSM. At this point one may wonder how with negligibly small higher FC one can have a screening regime at  $r > r_0$ . The answer was suggested in [14] where it was demonstrated that screening terms appear as an additive contribution to the Wilson loop which is not included in the cumulant expansion (2) and has a small coefficient  $O(N_c^{-2})$ . Therefore transition from the Casimir scaling regime to the screening regime occurs sharply at large  $T$  and  $r = r_0$  due to the definition of the static potential (1) and cannot be seen in the one-loop cumulant expansion in (2). For details the reader is referred to [4] and subsequent papers.

This work was partially supported by the joint grant RFFI-DFG, 96-02-00088G. The author is grateful to Dr. G.Bali for sending his lattice data and for correspondence, to A.M.Badalian, D.I.Diakonov, M.I.Polykarpov and G.t'Hooft for useful discussions, and V.I.Shevchenko for discussions and help in calculations.

- 
1. G.Bali, hep-lat/9908021.
  2. N.A.Campbell, I.H.Jorysz, and C.Michael, Phys. Lett. **B167**, 91 (1986); C.Michael, Nuc. Phys. (Proc. Suppl.) **B26**, 417 (1992); C.Michael, hep-ph/9809211; S.Deldar, hep-lat/9809137, hep-lat/9909077.
  3. H.G.Dosch, Phys. Lett. **B190**, 177 (1987); H.G.Dosch and Yu.A.Simonov, Phys. Lett. **B205**, 339 (1988); Yu.A.Simonov, Nucl. Phys. **B307**, 512 (1988), Yu.A.Simonov, Phys. Usp. **39**, 313 (1996).
  4. V.I.Shevchenko and Yu.A.Simonov, Phys. Rev. Lett. (subm.to).
  5. M.Peter, Phys. Rev. Lett. **78**, 602 (1997); Nucl. Phys. **B501**, 471 (1997); Y.Schroeder, Phys. Lett. **B447**, 321 (1999).

6. A.Di Giacomo and H.Panagopoulos, Phys. Lett. **B285**, 133 (1992); A.Di Giacomo, E.Meggiolaro, and H.Panagopoulos, Nucl. Phys. **B483**, 371 (1997); M.D'Elia, A.Di Giacomo, and E.Meggiolaro, Phys.Lett. **B408**, 315 (1997).
7. Yu.A.Simonov, Yad.Fiz. **54**, 192 (1991); H.G.Dosch, Progr. in Part Nucl. Phys. **33**, 121 (1994); O.Nachtmann, *Lecture Notes in Physics*, Springer, v.479, 1996, p.49; Yu.A.Simonov, hep-ph/9911237.
8. M.Faber, J.Greensite, and Š.Olejník, Phys. Rev. **D57**, 2603 (1998); hep-lat/9710039; S.Deldar, hep-ph/9912428.
9. L.Del Debbio, M.Faber, J.Greensite, and Š.Olejník, Phys. Rev. **D53**, 5891 (1996).
10. G.Poulis, Phys. Rev. **D54**, 6974 (1996); hep-lat/9608054.
11. E.M.Ilgenfritz, B.V.Martemyanov, S.V.Molodtsov et al., Phys. Rev. **D58**, 114508 (1998).
12. Yu.A.Simonov, Sov. J. Nucl. Phys. **50**, 310 (1989).
13. C.G.Callan, R.Dashen, and D.J.Gross, Phys. Lett. **B66**, 375 (1977); A.I.Veselov and M.I.Polikarpov, JETP Lett. **45**, 139 (1987); D.I.Diakonov, V.Yu.Petrov, and P.V.Pobylitsa, Phys. Lett. **B226**, 471 (1989); E.V.Shuryak, Nucl. Phys. **B328**, 85, 102 (1989).
14. J.Greensite and M.B.Halpern, Phys. Rev. **D27**, 2545 (1983).